

On ambiguity of definition of shear and spin-hall contributions to Λ polarization in heavy-ion collisions

Yu. B. Ivanov^{+*×1)}, A. A. Soldatov*

⁺Bogoliubov Laboratory for Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia

*National Research Nuclear University “MEPhI”, 115409 Moscow, Russia

×National Research Centre “Kurchatov Institute”, 123182 Moscow, Russia

Submitted 19 June 2022
Resubmitted 23 June 2022
Accepted 24 June 2022

DOI: 10.31857/S1234567822150010, EDN: jequnw

Non-central heavy-ion collisions at high energies are characterized by a huge global angular momentum of the order of 10^3 – $10^5 \hbar$, depending on the collision energy and centrality. Although a large part of the angular momentum is carried away by the spectator nucleons, its sizable fraction is accumulated in the created dense and highly excited matter, that implies a strong rotational motion of this matter. This matter is conventionally associated with a (participant) fluid because it is successfully described by the fluid dynamics. Such rotation leads to a strong vortical structure inside the produced fluid. Local fluid vorticity induces a preferential orientation of spins of emitted particles through spin-orbit coupling. The STAR Collaboration at the Relativistic Heavy-Ion Collider discovered the global polarization of emitted Λ hyperons, which indicated fluid vorticity of $\omega \approx (9 \pm 1) \times 10^{21} \text{ s}^{-1}$ [1]. This result exceeds the vorticity of all ever known fluids in nature. This discovery have opened an entirely new direction of research in heavy-ion physics.

The major part of applications to heavy-ion collisions was performed within thermodynamic approach in terms of hadronic degrees of freedom [2–4]. In the present paper, we discuss this thermodynamic approach. The key quantity of the thermodynamic approach is thermal vorticity

$$\varpi_{\mu\nu} = \frac{1}{2}(\partial_\nu \beta_\mu - \partial_\mu \beta_\nu), \quad (1)$$

where $\beta_\mu = u_\mu/T$, u_μ is collective local four-velocity of the matter, and T is local temperature. The corresponding mean spin vector of Λ particles with four-momentum p , produced around point x on freeze-out hypersurface is

$$S_{\varpi}^\mu(x, p) = -\frac{1}{8m}[1 - f(x, p)] \epsilon^{\mu\nu\alpha\beta} p_\nu \varpi_{\alpha\beta}(x), \quad (2)$$

where $f(x, p) = 1/\{\exp[(u_\nu p^\nu - \mu)/T] + 1\}$ is the Fermi-Dirac distribution function, m is mass of the Λ hyperon and μ is the baryon chemical potential.

It was recently realized [5–7] that there are other additional contributions to the mean spin vector, if the thermal equilibrium is *local*. These are the so-called thermal-shear (S_ξ^μ) and spin-Hall (S_ζ^μ) contributions:

$$S_\xi^\mu(x, p) = \frac{1}{4m}[1 - f(x, p)] \epsilon^{\mu\nu\alpha\beta} \frac{p_\nu n_\beta p^\rho}{(n \cdot p)} \xi_{\rho\alpha}, \quad (3)$$

$$S_\zeta^\mu(x, p) = \frac{1}{4m}[1 - f(x, p)] \epsilon^{\mu\nu\alpha\beta} \frac{p_\alpha n_\beta}{(n \cdot p)} \partial_\nu \zeta, \quad (4)$$

where $\zeta = \mu/T$,

$$\xi_{\mu\nu} = \frac{1}{2}(\partial_\mu \beta_\nu + \partial_\nu \beta_\mu), \quad (5)$$

is the thermal-shear tensor, and n is a four-vector that is the main subject of the discussion below. In [5], the n four-vector is defined as the time direction in the center-of-mass frame of colliding nuclei: $n_\beta = \hat{t}_\beta = (1, 0, 0, 0)$. Only the shear term was considered in [5]. In [6, 7], the n four-vector is identified with the four-velocity: $n_\beta = u_\beta$. In the mid-rapidity region these choices are very close because $u_\beta \approx \hat{t}_\beta$. However, at forward-backward rapidities, which are relevant to fixed-target polarization measurements [8–10], they may significantly differ.

In this paper, we consider consequences of these different choices at the example of the global polarization of Λ -hyperons. The global polarization is chosen because it allows a significant advance in the analytical treatment, if momentum acceptance is disregarded, and because it is relevant to fixed-target polarization measurements at moderately relativistic energies [8–10].

¹⁾e-mail: yivanov@theor.jinr.ru

The polarization of the Λ hyperon is measured in its rest frame, therefore the Λ polarization is

$$P^\mu(x, p) = S^{*\mu}(x, p)/S_\Lambda, \quad (6)$$

where $S_\Lambda = 1/2$ is the spin of the Λ hyperon, $S^\mu = S_\omega^\mu + S_\xi^\mu + S_\zeta^\mu$, and $S^{*\mu}$ is the mean Λ -spin vector in the Λ rest frame

$$\mathbf{S}^*(x, p) = \mathbf{S} - \frac{\hat{t} \cdot \mathbf{S}}{\hat{t} \cdot \mathbf{p} + m} \mathbf{p}_\Lambda \stackrel{\text{def}}{=} \mathbf{S} - \Delta \mathbf{S}, \quad (7)$$

where $\Delta \mathbf{S}$ is the boost correction. The zeroth component of $S^{*\mu}$ identically vanishes.

It is shown that alternative definitions of the thermal-shear contribution to the polarization in heavy-ion collisions, [5] on the one hand and [6, 7] on the other, result in very different corrections to the global Λ polarization averaged over wide range of momenta. The spin-Hall contribution to the polarization, defined accordingly to [6, 7], results in identically zero correction to the global Λ polarization, if averaged over all momenta of Λ 's. Only application of restrictive momentum acceptance and the boost (to Λ rest frame) correction result in nonzero global spin-Hall polarization. If the spin-Hall contribution were defined similarly to [5], the global spin-Hall polarization would be non-zero even without any acceptance and the boost correction.

This work was supported by MEPhI within the Federal Program "Priority-2030".

This is an excerpt of the article "On ambiguity of definition of shear and spin-hall contributions to polarization in heavy-ion collisions". Full text of the paper is published in JETP Letters journal. DOI: 10.1134/S0021364022601300.

1. L. Adamczyk, J.K. Adkins, G. Agakishiev et al. (STAR Collaboration), *Nature* **548**, 62 (2017); arXiv:1701.06657 [nucl-ex].
2. F. Becattini, V. Chandra, L. Del Zanna, and E. Grossi, *Annals Phys.* **338**, 32 (2013); arXiv:1303.3431 [nucl-th].
3. F. Becattini, I. Karpenko, M. Lisa, I. Upszal, and S. Voloshin, *Phys. Rev. C* **95**(5), 054902 (2017); arXiv:1610.02506 [nucl-th].
4. R.-h. Fang, L.-g. Pang, Q. Wang, and X.-n. Wang, *Phys. Rev. C* **94**(2), 024904 (2016); arXiv:1604.04036 [nucl-th].
5. F. Becattini, M. Buzzegoli, and A. Palermo, *Phys. Lett. B* **820**, 136519 (2021); arXiv:2103.10917 [nucl-th].
6. S.Y.F. Liu and Y. Yin, *JHEP* **07**, 188 (2021); arXiv:2103.09200 [hep-ph].
7. S.Y.F. Liu and Y. Yin, *Phys. Rev. D* **104**(5), 054043 (2021); arXiv:2006.12421 [nucl-th].
8. M.S. Abdallah, B.E. Aboona, J. Adam et al. (STAR), *Phys. Rev. C* **104**(6), L061901 (2021); arXiv:2108.00044 [nucl-ex].
9. K. Okubo for the STAR collaboration, *EPJ Web Conf.* **259**, 06003 (2022); arXiv:2108.10012 [nucl-ex].
10. R. Abou Yassine, J. Adamczewski-Musch, O. Arnold et al. (HADES Collaboration), arXiv:2205.15914 [nucl-ex]; F. J. Kornas, doi:10.26083/tuprints-00019763.