

ANYONS IN THE MAGNETIC FIELD AND FRACTIONAL QUANTUM HALL EFFECT

A.A. Ovchinnikov

Institute for Nuclear Research of the RAS,
117312, Moscow

Submitted 13 Januar 1992

We argue the incompressibility of a system of free anyons in the external magnetic field at special filling fractions. A new sequence of fractional quantum Hall effect states of the first level of hierarchy is proposed.

The characterization of all the possible incompressible states is one of the main goals of the complete theory of the fractional quantum Hall effect (FQHE). Besides the main states with the filling factors $\nu = 1/m$ (m is an odd integer) described by the Laughlin's variational wave function¹, a large number of other rational filling fractions are observed experimentally. At present time their classification is far from being completed. The first construction of the hierarchy of filling fractions^{2,3} was based on the quasiparticle wave functions of the Laughlin's type (or equivalently on the idea of condensation of the effective bosonic particles). The notion of binding of $m - 1$ flux quanta by electrons and integer quantum Hall effect of these composite objects was proposed by Jain⁴. The corresponding states are the lowest level of hierarchy of FQHE states. The field-theoretical realization of this scenario was proposed in ref.⁵ and ref.⁶. The Coulomb interaction is essential in the formation of these states. Recently the existence of a wider class of FQHE states of the first level of hierarchy was argued by Ma and Zhang⁷. It is known that the quasiparticles (quasiholes) in the basic $1/m$ FQHE states are fractionally charged and obey the fractional statistics. The latter can be understood either using the variational wave functions^{3,8} or in the framework of the description in terms of the bosonic variables⁹, where the quasiparticles correspond to the vortices which acquire the non-zero energy due to the Coulomb interaction. In ref.⁷ the incompressibility of the system of anyons in the magnetic field at special filling factors was claimed. If an integer number of the Landau levels in the effective magnetic field is filled, the system is incompressible even in the absence of the interaction, which is important in this scenario only for the formation of the quasiparticles. The arguments⁷ based on the perturbation theory in the small deviation from the Fermi-statistics are not convincing enough. Actually this deviation is not small. Besides the formation of the mean field cannot be described in the framework of perturbation theory. For instance the system of free anyons obeying the $1/N$ -statistics is compressible even for a large N .

In the present letter we propose the arguments in favor of the incompressibility of the system of anyons at special filling fractions using the description in terms of the Chern-Simons theory. We obtain a wider class of the filling fractions of FQHE states of the first level of hierarchy, which differs from that of ref.⁷ due to power of singular gauge transformation of the fermionic wave function into another

antisymmetric (fermionic) wave function. Our filling fractions are different from that found by Jain ⁴. The analysis of both quasiparticle and quasihole states is performed.

Consider the system of anyons (quasiparticles) with the statistical parameter α in respect to Bose statistics in the external magnetic field H_q (the charge of the anyons equals 1). The filling factor for anyons $\nu_a = 2\pi n_q / H_q$, where n_q is the density of anyons. Let us transform the multiparticle (Bose) wave function Φ according to $(z_i = x_i + iy_i, i, j = 1, \dots, N)$

$$\Phi(\vec{x}_1, \dots, \vec{x}_N) = \prod_{i < j} \left(\frac{z_i - z_j}{|z_i - z_j|} \right)^p \Psi(\vec{x}_1, \dots, \vec{x}_N), \quad (1)$$

where p - is an odd integer. The new wave function Ψ corresponds to Fermi-statistics. In the second-quantized form, introducing the gauge field a_μ , we obtain the Lagrangian:

$$L = i\psi^\dagger \partial_0 \psi + a_0(\psi^\dagger \psi - n_q) + \frac{1}{2M} \psi^\dagger (\partial - i\vec{a} - i\vec{A}_1)^2 \psi - \frac{1}{4\pi(\alpha - p)} \epsilon^{\mu\nu\alpha} a_\mu \partial_\nu a_\alpha, \quad (2)$$

where the field ψ obeys Fermi-statistics, \vec{A} is the vector potential corresponding to the homogeneous magnetic field of the magnitude $H_{eff} = H_q + 2\pi n_q(\alpha - p)$, and M is the anyon mass. The integer number of Landau levels in the effective magnetic field is filled at $\nu_a = (p + 1/n - \alpha)^{-1}$, where n is an integer, $n \neq 0$ (the sign of H_{eff} is not fixed). The fermion propagator has a gap equal to the energy difference between two Landau levels, the fermions can be integrated out in the action (2). In the one-loop effective action for the field a_μ the cancellation of the Chern-Simons term does not take place. Moreover the non-renormalization theorem ¹⁰ (for its application to the system of anyons see ref's ^{6,11}) leads to the conclusion that there are no perturbation theory corrections to the Chern-Simons term in the effective action ¹². That means that the collective excitation corresponding to the fluctuations of density (described by the field a_μ) has a gap (the field a_μ acquires a mass). The single-particle excitations are gapful due to the filling of the Landau levels. Thus for example the imaginary part of the correlator of two electromagnetic currents at small energies is equal to zero: the gapless excitations are absent. Thus the system at the filling factors considered is incompressible. It should be noted that the difference between the present case and the case of the Chern-Simons theory for the initial $1/m$ - state ^{5,6} is due to the degeneracy of the $\nu = 1/m$ - state in the absence of interaction. This degeneracy manifests in the existence of the vortices (quasiparticles) with zero creation energy ⁹. In our case these quasiclassical solutions are absent due to the non-integer value of $\alpha - p$ in the Lagrangian (2) which makes the perturbation theory (2) well defined even without the interaction. Alternatively, the exact wave function of the non-degenerate ground state can be constructed at $\nu_a = (p + 1/n - \alpha)^{-1}$ using the methods of ref. ¹³. This ground state corresponds to the classical configuration $a_\mu = 0$ in the effective action for the field a_μ . Thus, the long-wave fluctuations of this field are properly described by the effective action of the form ¹².

For the FQHE this scenario implies the following filling fractions. For the quasiparticles in the state with a given m we have $H_q = H/m$ and $\alpha = 1/m$.

The filling factor for the electrons is $\nu = (1/m)(1 + \nu_a/m)$. Thus we obtain the following quasiparticle states:

$$\nu^{qp} = \frac{1}{m - \frac{n}{np+1}}, \quad (3)$$

where n - is an integer ($n \neq 0$) and p - is an odd integer such that $n/(np + 1) > 0$ ($n > 0$). For the quasiholes we have $H_q = -H/m$ and $\alpha = 1/m$ and

$$\nu^{qh} = \frac{1}{m + \frac{n}{np-1}}, \quad (4)$$

where n - is an integer ($n \neq 0$) and p - is an odd integer such that $n/(np - 1) > 0$. Let us stress once more that the equations (3) and (4) represent the FQHE states of the first level of hierarchy. In fact our construction is in some sense the generalization of Jain's scheme to the first level of hierarchy. The same procedure can be performed both for quasiparticles of the next hierarchy levels and for the quasiparticle excitations of the Jain's states. At $p = 1$ the filling factors (3) and (4) coincide with the one of the two sequences of filling factors obtained by Jain (and with the sequence of states of ref.⁷). At $n = 1$ we get exactly the formulas of the first hierarchy level: the fermions at the completely filled single Landau level can be transformed into the bosons at zero effective magnetic field which corresponds to the construction of the hierarchy^{2,3}. As we have mentioned at $p = 1$ the states (3) and (4) coincide with the states of ref.⁴ (for $m = 3$: $(1/3), 2/5, 3/7, 4/9, 5/11, \dots$). At $p = 3$ the $m = 3$ sequence reads: $(1/3), 4/11, 7/19, \dots$ for quasiparticles and $(1/3), 2/7, 5/17, \dots$ for the quasiholes. For $p = 5$ the $m = 3$ sequences are $(1/3), 6/17, 11/31, \dots$ and $(1/3), 4/13, 9/29, \dots$ for the quasiparticles and the quasiholes respectively. For instance one can see that the fractions $4/11$ and $4/13$ that were observed experimentally are predicted. At the same time the existence of these states cannot be explained in the framework of the approach of ref.⁴. In general, the interpretation of the observed FQHE states as the states of high hierarchy level of the Haldane-Halperin hierarchical scheme is not quite natural. One can verify that all the FQHE states observed experimentally can be identified either with the states of the lowest level of hierarchy proposed in ref.⁴, or with the states given by our equations (3) and (4). As for the filling fractions predicted by both of these two schemes, much work is needed to determine the most adequate one. Clearly it depends on the dynamics of the formation of the quasiparticles i.e. on their size in comparison with the average distance between the quasiparticles.

-
1. R.B.Laughlin, Phys.Rev.Lett. **50**, 1395 (1983).
 2. F.D.M.Haldane, Phys.Rev.Lett. **51**, 605 (1983); R.B.Laughlin, Surface Science **141**, 1 (1984).
 3. B.I.Halperin, Phys.Rev.Lett. **52**, 1583 (1984).
 4. J.K.Jain, Phys.Rev.Lett. **63**, 199 (1989).
 5. A.Lopez and E.Fradkin, Phys.Rev. B **44**, 5246 (1991).
 6. A.A.Ovchinnikov, Pisma v ZhETF **54**, 579 (1991); Mod.Phys.Lett. A to appear.
 7. M.Ma and F.C.Zhang, Phys.Rev.Lett. **66**, 1769 (1991).

8. D.A.Arovas, R.Schreiffer and F.Wilczek, *Phys.Rev.Lett.* **53**, 722 (1984).
9. S.M.Girvin, In: *The quantum Hall effect*. Eds. R.E.Prange and S.M.Girvin, 1987; C.L.Kane, S.Kivelson and D.H.Lee, S.C.Zhang, *Phys.Rev.B* **43**, 3255 (1991); Z.F.Ezawa, M.Hotta and A.Iwazaki, Tohoku University preprint TU-376 (1991); *Phys.Rev. D* **44**, 452 (1991).
10. S.Coleman and B.Hill, *Phys.Lett. B* **159**, 184 (1985).
11. J.D.Lykken, J.Sonnenschein and N.Weiss, *Phys.Rev.D* **42**, 2161 (1990).
12. Y.H.Chen, F.Wilczek, E.Witten and B.I.Halperin, *Int.J.Mod.Phys. B* **3**, 1001 (1989); T.Banks and J.D.Lykken, *Nucl.Phys. B* **336**, 500 (1990); S.Randjbar-Daemi, A.Salam and J.Strathdee, *Nucl.Phys. B* **340**, 403 (1990).
13. M.Johnson and G.S.Girvin, *Phys.Rev. B* **41**, 6870 (1990); S.M.Girvin et. al., *Phys.Rev.Lett.* **65**, 1671 (1990); M.Greiter and M.Wilczek, preprint, 1991.