

ON THE FOUR-PHOTON DECAY OF THE NEUTRAL PION

E.L.Bratkovskaya, E.A.Kuraev, Z.K.Silagadze¹⁾

*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
141980 Dubna, Moscow Region, Russia*


Submitted 21 June 1995

The dominant contribution to the $\pi^0 \rightarrow 4\gamma$ branching ratio, coming from purely electromagnetic photon-splitting graph, is calculated. The result $Br(\pi^0 \rightarrow 4\gamma) \simeq (2.6 \pm 0.1) \cdot 10^{-11}$ is about three orders of magnitude below the present experimental limit.

Although a C-violating decay $\pi^0 \rightarrow 3\gamma$ is expected to have an extremely small branching ratio [1], beyond the reach of any present or future experimental facilities, its experimental study has attracted considerable attention [2] because any observed anomaly in this process would be a clear signal of a new physics.

Any $\pi^0 \rightarrow 3\gamma$ searching experiment has as a by-product, information about the allowed decay $\pi^0 \rightarrow 4\gamma$, that is a potential background for $\pi^0 \rightarrow 3\gamma$. The experimental upper limit on the branching ratio $Br(\pi^0 \rightarrow 4\gamma)$ was gradually improved [3-5] and lowered up to $2 \cdot 10^{-8}$ in [2]. Some theoretical estimates can be found in the literature for $Br(\pi^0 \rightarrow 4\gamma)$ [6-8], with rather broad ranges from 10^{-9} to 10^{-16} . In our opinion, the results of [8] are more reliable, the authors giving the most thorough investigation of the subject.

As argued in [8], the dominant contribution to the $\pi_0 \rightarrow 4\gamma$ branching ratio is expected to come from the purely electromagnetic photon-splitting graph of Figure, contributions from any hadronic intermediate states being less significant, especially for a PCAC-satisfying models. The calculation of this contribution is still absent to our best knowledge, and will be performed in the present note.



The electromagnetic photon-splitting graph

Using the standard covariant phase-space calculation technique [9] and factoring out some numerical constants from the decay amplitude, we can write

$$Br(\pi^0 \rightarrow 4\gamma) \simeq \frac{\Gamma(\pi^0 \rightarrow 4\gamma)}{\Gamma(\pi^0 \rightarrow 2\gamma)} = \frac{1}{6\pi} \left(\frac{\alpha}{8\pi}\right)^4 R, \quad (1)$$

¹⁾permanent address: Budker Institute of Nuclear Physics, 630090, Novosibirsk, Russia

where

$$R = \int_0^1 ds_1 \int_0^{s_1} ds_2 \int_{s_2/s_1}^{1-s_1+s_2} \frac{du_1}{\sqrt{\lambda(1, s_2, s'_2)}} \int_{u_2^-}^{u_2^+} du_2 \int_{-1}^1 \frac{d\zeta}{\sqrt{1-\zeta^2}} \cdot F(s_1, s_2, u_1, u_2, t_2(\zeta)). \quad (2)$$

In (2) we have introduced a dimensionless version of Kumar's invariant variables [9]:

$$\begin{aligned} s_1 &= \frac{1}{m^2}(q - k_1)^2, & s_2 &= \frac{1}{m^2}(q - k_1 - k_2)^2, \\ u_1 &= \frac{1}{m^2}(q - k_2)^2, & u_2 &= \frac{1}{m^2}(q - k_3)^2, \end{aligned} \quad (3)$$

m - being the pion mass.

One more invariant variable $t_2 = m^{-2}(q - k_2 - k_3)^2$ is a linear function of the integration variable ζ :

$$\begin{aligned} t_2 &= u_1 - \frac{1}{2}(1 + u_1)(1 - u_2) - \\ &- \frac{1}{2}(1 - u_1)(1 - u_2) \left[\xi\eta - \sqrt{(1 - \xi^2)(1 - \eta^2)}\zeta \right], \end{aligned} \quad (4)$$

where

$$\begin{aligned} \xi &= \frac{\lambda(1, s_2, s'_2) - (1 - s_1)^2 + (1 - u_1)^2}{2(1 - u_1)\sqrt{\lambda(1, s_2, s'_2)}}, \\ \eta &= \frac{(1 - s'_3)^2 - (1 - u_2)^2 - \lambda(1, s_2, s'_2)}{2(1 - u_2)\sqrt{\lambda(1, s_2, s'_2)}}, \end{aligned} \quad (5)$$

$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2(xy + xz + yz)$ is a conventional triangle function, and

$$s'_2 = 1 + s_2 - u_1 - s_1, \quad s'_3 = 2 - s_1 - u_1 - u_2. \quad (6)$$

At last, the limits of integration for the u_2 -variable in (2) are

$$u_2^\pm = 1 - \frac{1}{2}(u_1 + s_1) \pm \frac{1}{2}\sqrt{\lambda(1, s_2, s'_2)}. \quad (7)$$

Function F stands for a half sum of the squared helicity amplitudes:

$$F = \sum_{\{\lambda\}} |M_{+\lambda_2 \lambda_s \lambda_e}|^2. \quad (8)$$

To evaluate these helicity amplitudes, it is convenient to use the light-light scattering tensor from [10]. In fact, as virtuality of the intermediate photon is $\sim m$ and much bigger than the electron mass, we have used the asymptotic form of the light-light scattering amplitudes for the massless electron in the loop [11, 12].

Let us note, however, that we can not use the polarization vectors from [10] because of an additional photon and the need for photon permutations. Instead we have taken polarization vectors in the form which appeared useful in various QED calculations [13]:

$$\begin{aligned} \varepsilon_{\mu}^{(\lambda_m)}(k) &= N_m \left[q^{(m)} \cdot k p_{\mu}^{(m)} - p^{(m)} \cdot k q_{\mu}^{(m)} + i \lambda_m \varepsilon_{\mu\nu\lambda\sigma} p^{(m)\nu} k^{\lambda} q^{(m)\sigma} \right], \\ N_m^{-1} &= 2\sqrt{p^{(m)} \cdot q^{(m)} p^{(m)} \cdot k q^{(m)} \cdot k}, \quad m = 1 \div 4, \quad p^{(m)2} = q^{(m)2} = 0. \end{aligned} \quad (9)$$

Where for various photons we take

$$\begin{aligned} p^{(1)} &= k_2, & p^{(2)} &= k_3, & p^{(3)} &= k_4, & p^{(4)} &= k_1, \\ q^{(1)} &= k_4, & q^{(2)} &= k_1, & q^{(3)} &= k_2, & q^{(4)} &= k_3. \end{aligned} \quad (10)$$

The polarization vectors from [10] can also be expressed in this form ($8\Delta = k_2 \cdot k_3 k_3 \cdot k_4 k_2 \cdot k_4$):

$$\begin{aligned} u_{\mu}^{(-\lambda_2)} &= \frac{1}{4\sqrt{2\Delta}} [k_4 \cdot k_2 k_{3\mu} - k_3 \cdot k_2 k_{4\mu} + i \lambda_2 \varepsilon_{\mu\nu\lambda\sigma} k_3^{\nu} k_2^{\lambda} k_4^{\sigma}], \\ u_{\mu}^{(\lambda_3)} &= \frac{1}{4\sqrt{2\Delta}} [k_2 \cdot k_3 k_{4\mu} - k_4 \cdot k_3 k_{2\mu} + i \lambda_3 \varepsilon_{\mu\nu\lambda\sigma} k_4^{\nu} k_3^{\lambda} k_2^{\sigma}] \equiv \varepsilon_{\mu}^{(\lambda_3)}, \\ u_{\mu}^{(-\lambda_4)} &= \frac{1}{4\sqrt{2\Delta}} [k_3 \cdot k_4 k_{2\mu} - k_2 \cdot k_4 k_{3\mu} + i \lambda_4 \varepsilon_{\mu\nu\lambda\sigma} k_2^{\nu} k_4^{\lambda} k_3^{\sigma}]. \end{aligned} \quad (11)$$

But $u_{\mu}^{(-\lambda_2)}$ and $u_{\mu}^{(-\lambda_4)}$ differ by the phase factors from $\varepsilon_{\mu}^{(\lambda_2)}$ and $\varepsilon_{\mu}^{(\lambda_4)}$ (note that, in contrast with [10], $u_{\mu}^{(-\lambda_2)}$ corresponds to the $+\lambda_2$ circular polarization for the second photon because now it is also outgoing). Therefore, while using the expressions from [10], we should not forget the relevant phase factors. For example,

$$u_{\mu}^{(\lambda_2)} \cdot \varepsilon^{(\lambda_2)} = \frac{N_2 k_2 \cdot k_3}{2\sqrt{2\Delta}} \Phi(\lambda_2; 1234),$$

where

$$\begin{aligned} \Phi(\lambda; 1234) &= k_1 \cdot k_2 k_3 \cdot k_4 + k_1 \cdot k_3 k_2 \cdot k_4 - k_1 \cdot k_4 k_2 \cdot k_3 + \\ &+ i \lambda [k_1, k_2, k_3, k_4], \\ [k_1, k_2, k_3, k_4] &= \varepsilon_{\mu\nu\lambda\sigma} k_1^{\mu} k_2^{\nu} k_3^{\lambda} k_4^{\sigma}. \end{aligned} \quad (12)$$

Owing to the remarkable cyclic symmetry in the definition (9), (10) of the $\varepsilon_{\mu}^{(\lambda_i)}$ polarization vectors, for the $\pi^0 \rightarrow 4\gamma$ helicity amplitudes we get (remind that some numerical factors have already been taken out in (1)):

$$\begin{aligned} M_{\lambda_1 \lambda_2 \lambda_3 \lambda_4} &= \frac{1}{k_1 \cdot k_2 k_1 \cdot k_3 k_1 \cdot k_4 k_2 \cdot k_3 k_2 \cdot k_4 k_3 \cdot k_4} \cdot \\ &\cdot \sum_{cyclic} \frac{\Phi(\lambda_2; 1234) \Phi(\lambda_4; 1432)}{k_2 \cdot k_4} \left\{ A(\lambda_1; 1234) \varepsilon_{-\lambda_2, \lambda_3, \lambda_4}^{(1)}(234) + \right. \\ &+ \left. B(\lambda_1; 1234) \varepsilon_{-\lambda_2, \lambda_4, \lambda_3}^{(1)}(243) + 2C(\lambda_1; 1234) \varepsilon_{-\lambda_2, \lambda_3, \lambda_4}^{(2)}(234) \right\}. \end{aligned} \quad (13)$$

Here the summation extends over simultaneous cyclic permutations of $(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$ and (k_1, k_2, k_3, k_4) :

$$\sum_{\text{cyclic}} F(1234) = F(1234) + F(2341) + F(3412) + F(4123).$$

The $\epsilon_{\{\lambda\}}^{(1)}$ and $\epsilon_{\{\lambda\}}^{(2)}$ amplitudes have been defined in [11] (see also [12]). For the A, B and C functions we have

$$\begin{aligned} A(\lambda_1; 1234) &= \epsilon^{\mu\nu\lambda\sigma} k_\mu^1 q_\nu \epsilon_\lambda^{(\lambda_1)} \left[k_3 - \frac{(q-k_1) \cdot k_3}{(q-k_1) \cdot k_2} k_2 \right]_\sigma, \\ B(\lambda_1; 1234) &= -\epsilon^{\mu\nu\lambda\sigma} k_\mu^1 q_\nu \epsilon_\lambda^{(\lambda_1)} \left[k_4 - \frac{(q-k_1) \cdot k_4}{(q-k_1) \cdot k_2} k_2 \right]_\sigma, \\ C(\lambda_1; 1234) &= i \epsilon^{\mu\nu\lambda\sigma} k_\mu^1 q_\nu \epsilon_\lambda^{(\lambda_1)} \epsilon_{\sigma\mu'\nu'\lambda'} (k_1 - q)^{\mu'} k_2^{\nu'} k_3^{\lambda'}. \end{aligned} \quad (14)$$

The polarization vectors (9) and (11) become ill-defined for collinear photons. Fortunately, this kinematical region gives a negligible contribution to the decay width. In fact the corresponding fictitious kinematical singularities don't cause any considerable trouble in numerical calculations because the phase factors also vanish for collinear photons and only the singularities corresponding to the three simultaneously collinear photons remain.

There are no infrared divergencies in our problem (when the energy of any photon goes to zero), as is easily seen from the explicit expressions of the $\epsilon^{(1)}, \epsilon^{(2)}$ amplitudes. The contribution of the muon in the fermion loop can be neglected at least by a factor $(\omega/m_\mu)^3$, $\omega \sim m/4$, ω is the mean photon energy, due to the known low energy behavior of the light-light scattering amplitude.

The numerical calculations give the result

$$Br(\pi^0 \rightarrow 4\gamma) \simeq (2.6 \pm 0.1) \cdot 10^{-11}. \quad (15)$$

This is about three orders of magnitude below the present experimental limit.

-
1. D.A.Dicus, Phys. Rev. **D12**, 2133 (1975).
 2. J.McDonough, R.D.Bolton, D.P.Grosnick et al., Phys. Rev. **D38**, 2121 (1988).
 3. L.B.Auerbach, V.L.Highland, R.J.Macek et al., Phys. Lett. **78B**, 353 (1978).
 4. L.B.Auerbach, V.L.Highland, R.J.Macek et al., Phys. Lett. **90B**, 317 (1980).
 5. V.N.Bolotov, S.N.Gninenko, R.M.Dzhilkibaeb et al., Pis'ma Zh. Eksp. Teor. Fiz. **43**, 405 (1986).
 6. A.V.Tarasov, Yad. Fiz. **5**, 626 (1967).
 7. D.Parashar and P.N.Dobson, Jr., Phys. Rev. **D12**, 77 (1975).
 8. R.L.Schult and B.-L.Young, Phys. Rev. **D6**, 1988 (1972).
 9. R.Kumar, Phys. Rev. **185**, 1865 (1969); E.Byckling and K. Kajantie, *Particle Kinematics*, Wiley, London, 1973.
 10. V.Constantini, B.De Tollis, and G.Pistoni, *Novo Cim.* **2A**, 733 (1971).
 11. V.N.Ivanchenko, E.A.Kuraev, and V.S.Fadin, Preprint INP-79-13, Novosibirsk, 1979.
 12. V.Baier, E.Kuraev, and V.Fadin, *Sov. J. Nucl. Phys.* **31**, 364 (1980); M.L.Laursen, K.O.Mikaelian, and M.A.Samuel, Phys. Rev. **D23**, 2795 (1981); M.L.Laursen and M.A.Samuel, *Z. Phys.* **C14**, 325 (1982).
 13. P.de Causmaecker, R.Gastmans, W.Troost, and T.T.Wu, *Nucl. Phys.* **B206**, 53 (1982).